شركة الحديثـة لصناعة الأدوية البيطريـة و المبيـدات الزراعيـة

The Modern Company for Veterinary Medicines & Agricultural Pesticides Industries

Pharmaceutical

POVIDONE - MOFED

Cow Camel

Goat Lamb Horse

Cat Dog

Ointment - Used in the treatment or prevention of local infections

INDICATIONS:

As an aid in the treatment or prevention of local infections in cases of wounds, abscesses, burns and fungal infections, e.g. disinfection of the navel, removal of horns, castration, ringworm, disinfection prior to injection or surgery.

DOSAGE AND ADMINISTRATION:

After cleaning the infected area with warm water and soap, dry thoroughly, and then apply the ointment daily until healing. Apply ointment to the wound and surrounding area. Usually one treatment is sufficient to prevent infection. If used therapeutically when infection is present, treatment may be repeated several times daily as required. May be bandaged. In case of deep or puncture wounds, or serious burns consult a veterinarian. Do not use if hypersensitivity to iodine is suspected. If irritation, redness or swelling develops, discontinue treatment. Avoid application over large skin areas.

COMPOSITION:

Each 1 gm contains:

Povidone Iodine	100 mg
Vaseline add to	1 gm

WITHDRAWAL PERIOD:

None

200 gm

(below 30°C)

شركة الحديثــة لصناعة الأدوية البيطريــة و المبيــدات الزراعيــة
The Modern Company for Veterinary Medicines & Agricultural Pesticides Industries

Pharmaceutical

بوفيدون - موفيد

Cow Camel
Goat Lamb Horse
Cat Dog

مرهم - يستخدم في علاج أو الوقاية من الالتهابات الموضعية

التركيبة :

كـل ١ غم يحتـوي :

بوفيدون اليود ١٠٠ ملغم
فازلين حتى ١ غم

فترة السحب :

لايوجد

٢٠٠ غم

(أقل من ٣٠ درجة مئوية)

الخواص العلاجية والإستطبابات:

يستعمل كعامل مساعد في علاج أو الوقاية من الالتهابات الموضعية في حالات الجروح، الخرّاجات، الحروق والالتهابات الفطرية مثل: تعقيم السرة، الأذان، إزالة القرون، الاخصاء ، القوباء الحلقية ، التعقيم قبل الحقن أو العمليات الجراحية.

الجرعات وطريقة الاستعمال :

بعد تنظيف المنطقة المصابة بالماء الدافئ والصابون، تجفف جيداً، ثم يُدهن المرهم يومياً حتى الشفاء التام.
يدهن المرهم على الجرح والمنطقة المحيطة به. عادةً ما تكون المعالجة لمرة واحدة كافية للوقاية من العدوى.
وإذا استُخدم علاجياً عند وجود التهاب، فقد يتطلب تكرار المعالجة عدة مرات يومياً حسب الحاجة.
يمكن تضميد المنطقة المصابة.
في حالة الجروح العميقة أو الجروح الناتجة عن وخز أو الحروق الشديدة، يجب استشارة الطبيب البيطري.
لا يُستخدم الدواء إذا كان هناك شك بوجود حساسية تجاه مادة اليود.
عند حدوث تهيج أو احمرار أو تورم، يجب وقف العلاج فوراً.
يُمنع استخدام المرهم على مساحات كبيرة من الجلد.