



شركة الحديثة لصناعة الأدوية البيطرية و المبيدات الزراعية
The Modern Company for Veterinary Medicines & Agricultural Pesticides Industries

Feed Additives

Move Electrolyte Plus Powder



Cow



Poultry



Goat



Lamb

Promotes Health and Growth in Birds
Increases Resistance Against Dehydration and Heat Stress

INDICATIONS:

- It is used for chicks in day one of reception to increase their activity after transportation stress and exposure to cold or heat.
- It is used to prevent dehydration caused by pathogenic diarrhea cases.
- Given in high temperatures causing sudden death due to heat retention (in this case the dose is doubled)
- Helps to activate liver functions and kidney detoxification.

DOSAGE AND ADMINISTRATION:

1 gm/L of drinking water for 3-5 days

COMPOSITION:

Each 1 kg contains:

Probiotic 10,000,000,000 cell, vitamin C 15,000 mg, vitamin K3 10,000 mg, vitamin E acetate 10,000 I.U, Sorbitol 60,000 mg, Sodium citrate 65,000 mg, Potassium chloride 50,000 mg, Calcium lactate 17,000 mg, Sodium chloride 8,000 mg, Sodium bicarbonate 30,000 mg, Sodium phosphate 8,000 mg, Magnesium sulfate 20,000 mg, DL-methionine 20,000 mg, L-lysine 10,000 mg, Amino Acids (lysine, alanine, arginine, aspartic acid, cystine, glutamic acid, glycine, histidine, L-leucine, L-isoleucine, phenylalanine, proline, serine, threonine, tyrosine) 50,000 mg.

WITHDRAWAL PERIOD:

None



1000 gm



(below 30°C)

