شركة الحديثـة لصناعة الأدوية البيطريـة و المبيـدات الزراعيـة

The Modern Company for Veterinary Medicines & Agricultural Pesticides Industries

Pharmaceutical

NAVOMOVED PLUS

> Cow Poultry

> Goat Lamb

water soluble powder

INDICATIONS:

Navomoved is a synergistic combination helps to stimulate kidney and liver functions. It reduces high temperature in bird caused by bacteria and viral pathogens especially in case (New Castle, Gumboro, E. coli....).
Navomoved lowers body temperature.
Navomoved is used in severe stress situations which cause weight loss and production such as (cutting beaks - transport - weight loss - oil and watery vaccines, cold and congestion ... etc).
Helps to increase the degree of food conversion and strengthens the egg shell and helps to modify lack of egg production caused by vitamin E deficiency.

DOSAGE AND ADMINISTRATION:

Poultry: 1 g per 1-2 liters of drinking water for 3 days

COMPOSITION:

Each 1 gm contains:

NOVELGEN 350 mg
Vitamin C 100 mg
Vitamin K3 20 mg
Vitamin E 30 mg
Sorbitol 10 mg
Potassium Citrate 10 mg

WITHDRAWAL PERIOD:

4 days

500 gm

(below 30°C)

Pharmaceutical

شركة الحديثــة لصناعة الأدوية البيطريــة و المبيــدات الزراعيــة

The Modern Company for Veterinary Medicines & Agricultural Pesticides Industries

نـافوموفيد بلص

Cow Poultry

Goat Lamb

مسحوق ذواب في الماء

التركيبة :

كل ١ غم يحتوي على:

نوفلجين ٣٥٠ ملغم
فيتامين سي ١٠٠ ملغم
فيتامين ك٣ ٢٠ ملغم
فيتامين هـ ٣٠ ملغم
سوربيتول ١٠ ملغم
سترات البوتاسيوم ١٠ ملغم

فترة السحب :

٤ أيام

٥٠٠ غم

(أقل من ٣٠ درجة مئوية)

الخواص العلاجية والإستطبابات:

نافوموفيد تركيبة فعالة تعمل على تنشيط وظائف الكلية والكبد، تخفض حرارة جسم الطائر الناتجة عن الإصابات البكتيرية والفيروسية التي تؤدي إلى رفع حرارة الجسم وخاصة في حالات نيوكاسل، الجمبورو، والإيكولاي....
نافوموفيد يخفض حرارة الجسم.
نافوموفيد يستخدم في حالات الإجهاد الشديد التي تسبب انخفاض الوزن والإنتاج مثل:
قص المنقار - النقل - انخفاض الوزن - اللقاحات الزيتية والمائية - البرد والازدحام .... الخ.
يساعد في رفع كفاءة التحويل الغذائي ويقوي قشرة البيض ويساهم في تعديل نقص الإنتاج الناتج عن نقص فيتامين هـ.

الجرعات وطريقة الاستعمال :

الدواجن: ١ غرام لكل ١-٢ لتر ماء شرب لمدة ٣ أيام