شركة الحديثـة لصناعة الأدوية البيطريـة و المبيـدات الزراعيـة

The Modern Company for Veterinary Medicines & Agricultural Pesticides Industries

NEW

Pharmaceutical

Ampicillin Move Plus

> Cow Poultry

> Goat Lamb

Water Soluble Powder - For GIT and Systemic Infections

INDICATIONS:

Ampicillin Move Plus is indicated for the treatment of infections caused by E.coli, Clostridium, Salmonella, Brucella, Proteus, Klebsiella, Streptococci, Staphylococci, Corynebacterium, Haemophilus, Pasteurella, and Shigella in different animals.
Ampicillin Move Plus is used in cases of Bacterial enteritis and Necrotic enteritis in poultry.
It is also indicated for Pneumonia, Septicemia, Diarrhea and Enteritis in calves and lambs

DOSAGE AND ADMINISTRATION:

Poultry: 1 gm per 2 liters of drinking water for 3-5 days
Calves, lambs and other animals: 1-2 gm per 10 kgs of body weight via drinking water for 3-5 days, the dose can be doubled in severe cases.

COMPOSITION:

Each 1 gram contains:

Ampicillin Trihydrate	200 mg
Colistin Sulfate	800.000 I.U

WITHDRAWAL PERIOD:

Meat: 7 days
Eggs: 3 days

500 gm

(below 30° C)

شركة الحديثــة لصناعة الأدوية البيطريــة و المبيــدات الزراعيــة
The Modern Company for Veterinary Medicines & Agricultural Pesticides Industries

Pharmaceutical

جديد

أمبيسيلين موفي بلص

Cow Poultry

Goat Lamb

بودرة قابلة للذوبان في الماء - لعلاج التهابات الجهاز الهضمي والجهازي

التركيبة :

يحتوي كل جرام على:
أمبيسلين تراي هيدرات ٢٠٠ مجم
كوليستين سلفات ٨٠٠,٠٠٠ وحدة دولية

فترة السحب :

اللحوم: ٧ أيام
البيض: ٣ أيام

٥٠٠ جم

(أقل من ٣٠ درجة مئوية)

الخواص العلاجية والإستطبابات:

يستخدم أمبيسيلين موفي بلص لعلاج الالتهابات التي تسببها البكتيريا القولونية، والكلوستريديوم، والسالمونيلا، والبروسيلا، والبروتيوس، والكلبسيلا، والعقديات، والمكورات العنقودية، والكورينباكتيريوم، والهايموفيلوس، والباستوريلا، والشيجيلا في الحيوانات المختلفة.
يستخدم أمبيسيلين موفي بلص في حالات التهاب الأمعاء البكتيري والتهاب الأمعاء النخرية في الدواجن.
كما يستخدم لعلاج الالتهاب الرئوي، وتسمم الدم، والإسهال والتهاب الأمعاء في العجول والحملان

الجرعات وطريقة الاستعمال :

الدواجن: ١ جرام لكل ٢ لتر من مياه الشرب لمدة ٣-٥ أيام
العجول والحملان والحيوانات الأخرى: ١-٢ جرام لكل ١٠ كجم من وزن الجسم عن طريق مياه الشرب لمدة ٣-٥ أيام، ويمكن مضاعفة الجرعة في الحالات الشديدة.